KEYLINA, Kh.V.

Climacteric hemorrhages. Zdrav.Bel. no.3:15-16 '62.

(MIRA 15:5)

1. Iz Baranovichskogo gorodskogo rodil'nogo doma (glavnyy vrach zasluzhennyy vrach BSSR V.P. Shchegoleva).

(HEMORRHAGE, UTERINE) (CLIMACTERIC)

KEYLINA, R. Ya.

"Biochemical Processes in the Brain in Birect Irradiation by X-rays,"
N. N. Blokhin, B. M. Grayevskaya, and R. Ya. Keylina (Cent. Roentgenob.
Radiol. Cancer Inst., Leningrad.) Byull. Eksptl. Biol. Med., 23, 338-42 (1947)

X-ray irradiation of dog brain, at 160 kv., at 23 mm. distance by using 0.5 Cu-3.0 Al filter, each dog receiving a total of 5 unit skin doses of irradiation in 4 exposures with alternate irradiation of the right and the left temple area, was investigated in respect to biodhem. effects by detn. of blood sugar (femoral artery and sinus venosus cerebri), spinal fluid sugar, blood serum protein, and the albumin-globulin fractions of the latter. As the total irradiation increased, the total serum portein rose until on the 90th day (after the 1st irradiation) it received 200% of the initial value. In the same period the albumin fractions rose by only 26% of arterial and 32% of venous blood, while the globulin fraction rose 324 and 323%, resp. The abs. amt. of protein in the spinal fluid remained within exptl. variations, but its albumin/globulin ration gradually changed to 1.0 from 0.56. Arterial blood sugar remained normal, but it fell in the venous blood, until the sugar utilization by t the brain at the 90-day period was 370% of initial.

Keylina R. Ya

BLOKHIN, N.N.; GRAYEVSKAYA, B.M.; KEYLINA, R.Ya.

Biochemical functional test in certain forms of hypertension.

Vop.med.khim. 3:52-57 '51. (MIRA 11:4)

1. Otdel biokhimii TSentral'nogo rentgenologicheskogo, radiologicheskogo i rakovogo instituta, Leningrad.
(HYPERTENSION)

KSYLLIA, R. Ya.

JSSK/Medicine - Radiation Effects

Nov/Dec 53

"Effects of Aneurin (Vitamin B₁) on the Glycogen Formation Function of Liver of Rats During General Exposure to Various Doses of X-Rays," B. M. Grayevskays, R. Ya. Keylina, and S. E. Manoylov, Dept of Blochem, Central Roentgenologic and Cancer Inst, Min of Health USSR

Vest Rentgen 1 Radiol, No 6, pp 22-25


Expts conducted on 15 rats showed that the glycogen formation function of the liver may be partially restored if vitamin B is given after rats have


275131

been exposed to X-rays. Enough evidence was collected to justify the assumption that the quality of biochem changes in the organism of rats depends upon the extent of exposure to X-rays: small doses (500 r) of X-rays suppress the oxidation systems of the whole organism: larger doses (1,000 and 2,000 r also cause breakdown of the substances that oxidize. It is possible that interference with the synthesis of the dehydrogenase coenzyme takes place and that vitamin B loses its capacity to combine with phosphoric acid. This follows from a reduction of the restorative effect of vitamin B on the


"APPROVED FOR RELEASE: 09/17/2001

CIA-RDP86-00513R000721620005-8





				: ·				·
The state of the s	pada tankan pada nagas settakan di tang basak set sa sebangan	and the second s	and the second second second second second		*.	*************	fdhuib is abh Aprila (bir sí s	o enqu
	VA	tion of fonizing	radiation on certain	phases of carbo-	۰. ۵			
	VAc hydr and (195	ction of fonizing rate exchange in R. Ya. Kellina. 5),-A review wi	radiation on certain animal organism. Uspekhi Sovressens th 67 references.	n phases of carbo- B. M. Graevskaya P of Piol. 40, "30-8	10 2			
	VAc hydr and (195	ction of ionizing rate exchange in R. Ya. Kelling, 5)A review wi	radiàtion on certain animal organism. I Uspekhi Sovressenn ith 67 references.	n phases of carbo- B. M. Graevskaya I set Riol. 40, "30-8 J. A. Stekol	10 2			
	VAC hydr and (195	ction of ionizing ate exchange in R. Ya. Keilina. 5)A review wi	radiation on certain animal organism. I Uspekhi Sovressenn th 67 references.	n phases of carbo- B. M. Graevskaya I tol Biol. 40, "30-8 J. A. Stekol	10 2			
	VAc hydr and (195	ction of ionizing rate exchange in R. Ya. Kellina. 5)A review wi	radiation on certain animal organism. I Uspekhi Sovressenn th 67 references.	n phases of carbo- B. M. Graevskaya of Biol. 40, "30-8 J. A. Stekol	10 2	-		
	VAc hydr and (195	ction of ionizing ate exchange in R. Ya. Keilina. S A review wi	radiation on certain animal organism. I Uspekhi Sovressens th 67 references.	n phases of carbo- R. M. Graevskaya P of Biol. 40, 30-8 J. A. Stekol	10 2		·	
	VA hydr and (195	ction of ionizing rate exchange in R. Ya. Keilina. 5)A review wi	radiation on certain animal organism. Uspekhi Sovresenz th 67 references.	n phases of carbo- R, M. Graevskaya P of Riol. 40, 30-8 J. A. Stekol	¹⁰ ②			
	VA hydr and (195	ction of ionizing rate exchange in R. Ya. Keilina, 55,A review wi	radiation on certain animal organism. Uspekhi Sovressen th 67 references.	n phases of carbo- R. M. Graevskaya P od Biol. 40, 30-8 J. A. Stekol	¹⁰ ②	-		



KEYLINA, R. YA.

"Changes of the Sugar Content of Blood Leaving the Brain and Muscles After Total Irradiation of Dogs by X Rays," by R. Ya. Keylina, Vopr. Radiobiologii (Problems of Radiobiology), Leningrad, 1956, pp 347 - 351 (from Referativnyy Zhurnal -- Khimiya, Biologicheskaya Khimiya, No 1, 10 Jan 57, Abstract No 702

After irradiation of dogs by X rays (300 - 500 r), the sugar consumption of the brain sharply decreased, reaching zero toward the 40th day, but starting with the 50th day there was an insignificant release of sugar into the circulating blood. However, the muscles at this time started to use up sugar intensely.

Following total irradiation of dogs by lethal doses (1,000 r), a high sugar content was maintained for 4 hours in the arterial blood subsequent to adrenalin injections.

Shifts detected in the carbohydrate metabolism following X-ray irradiation are evidences of disturbances of a number of enzyme systems participating in the breakdown and oxidation of glucose.

Sum. 1305


APPROVED FOR RELEASE: 09/17/2001 CIA-RDP86-00513R000721620005-8"

"Change of Carbohydrate Metabolism Due to Total Irradiation of an Animal Organism by X Rays," by B. M. Grayevskaya and R. Ya. Keymina, Voprosy Radiobiologii (Problems of Radiobiology), Leningrad, 1956, pp 352-356 (from Referativnyy Zhurnal -- Khimiya, Biologicheskaya Khimiya, No 24, 25 Dec 56, p 80, Abstract No 23449)

Subcutaneous injections of adrenalin to dogs who were subjected to total irradiations by 300 to 500 r did not cause increased blood glucose content. Liver glycogen content decreased 80% in 48 hours after irradiaation.

Glucose administration caused no glycogen synthesis in the liver of rats who had been subjected to 500, 1,000, or 2,000 r.

Sum 0.37



"APPROVED FOR RELEASE: 09/17/2001

CIA-RDP86-00513R000721620005-8

TATARSKIY, M.B.; KEYLINA, R.Ya., kond.biol.neuk, neuchnyy red.; VIADIMIRSKIY, D.M., ted.iva-va; Gurdžhiyeva, A.M., tekhn.red.

[Tagged atoms end their role in the understanding of vital phenomena]

| Metod mechenyth atomov i ego rol' v poznanii zhiznennykh iavleuii.
| Leningrad, 1957. 52 p. (MIRA 11:5)

(Redioactive tracers)

T-13 USSR / Human and Animal Physiology. Effect of Physical Factors.

: Ref Zhur - Biologiya, No 1, 1959, No. 3968 Abs Jour

Author

: Grayevskaya, B. M.; Keylira, R. Ya.

Inst

: Not given

Title

: The Decrease of Sensitivity of Animals to the Effect of Roentgen Rays in a Lethal Dose by Their Preliminary

Irradiation with Non-Lethal Doses

Orig Pub

: Tr. Vses. konferentsii po med. radiol. Eksperim. med.

radiol. M., Medgiz, 1957, 183-185

Rlas put. in BiofiziKA 1, No. 3, 232-236, 1956

Abstract

: The survival of animals subjected to a total irradiation of 1000 r (lethal dose) constituted respectively 50 and 100% 150 days (34 rats) and 120 - 150 days (8 dogs) after preliminary irradiation with a dose of 500 r. Repeated action, as differentiated from a single action in the same dose, was accompanied by a loss severe symptomatology and did not lead to a decrease of Hb content

Card 1/2 Cent. Sei Les. Reentgens. Ladislogs Inst.

127 Min Weelth, 555R, Lenerged

DETOCKA SERBENDA MENDENDENDE SERBENDENDENDENDENDE MERENDENDE DE SE

MANOYLOV, S.Ye.; GRAYEVSKAYA, B.M.; KEYLINA, R.Ya.

Use of some biological preparations (campolonum, vitamin B₁ and adenosinetriphosphoric acid) as prophylactic and therapeutic drugs in radiation sickness. Vop.radiobiol. 2:426-430 '57. (MIRA 12:6)

1. Sotrudniki "Sentral'nogo nauchno-issledovatel'skogo rentgenoradiologicheskogo instituta Ministerstva zdravookhraneniya SSSR. (VITAMINS-B) (RADIATION SICKMESS) (ADENOSIMETRIPHOSPHORIC ACID)

"HERT'S COMMENT DESIGNATION OF THE STATE OF

KEYLINA, R.Ya.

State of carbohydrate metabolism during the use of certain substances in the prophylaxis of radiation sickness (insulin, adrenalin, vitamin B, adenosinetriphosphoric acid, etc.).

Vop.radiobiol. 2:437-444 '57. (MIRA 12:6)

1. Sotrudnik TSentral'nogo nauchno-issledovatel'skogo rentgenoradiologicheskogo instituta Kinisterstva zdravookhraneniya SSSR. (RADIATION PROTECTION) (CARBOHYDRATE METABOLISM)

APPROVED FOR RELEASE: 09/17/2001 CIA-RDP86-00513R000721620005-8"

GRAYEVSKAYA, B.M., KEYLINA R.YA.

With regard to I.G. Krasnykh and S.P. IArmonenko's letter published in "Biofizika". Biofizika 3 no.4:527-528 58 (MIRA 11:8) (X RAYS--PHYSIOLOGICAL EFFECT)

KEYLINA, R.Ya.: KOMAROV, Ye.I.

Reflex mechanisms of changes in blood sugar level associated with local irradiation of the small intestine by radioactive strontium [with summary in English]. Fiziol.zhur. 44 no.2:148-152 F '58.

(MIRA 11:5)

1. TSentral'nyy nauchno-issledovatel'skiy rentgeno-radiologicheskiy institut, Leningrad.

(INTESTINES, SMALL, eff. of radiations on radiostrontium irradiation of exteriorized small intestine in dog, eff. on blood sugar levels, reflex mechanisms (Rus) (STRONTIUM, radioactive

irradiation of exteriorized small intestine of dog, eff. on blood sugar levels, reflex mechanisms (Rus)

(BLOOD SUGAR, physiol, eff. of radiostrontium irradiation of exteriorized small intestine of dog, reflex mechanisms (Rus)

GRAYEVSKAYA, B.M.; KEYLINA, R.Ya.

Role of the adrenals in certain metabolic disorders in irradiated organisms. Med. rad. 4 no.3:21-25 Mr '59. (MIRA 12:7)

1. Iz otdela biokhimii (zav. - prof. S.Ye. Manoylov) Tšentral'nogo nauchno-issledovatel'skogo rentgeno-radiologicheskogo instituta Ministerstva zdravookhraneniya SSSR.

(ROENTGEN RAYS, effects,
on carbohydrate & protein metab., eff. of adrenalectomy
(Rus))
(CARBOHYDRATE, metab.
eff. of x-rays in adrenalectomized animals (Rus))
(PROTEIN, metab.
same)
(ADRENALECTOMY, effects,
on carbohydrate & protein metab. responses to x-rays
(Rus))

THE WASHERS THE PROPERTY OF TH

KEYLINA, R.Ya.

Effect of whole body X-irradiation on the glycogen content of the rat liver. Biokhimia 24 no.6:966-999 N-D 159.

(MIRA 13:5)

1. Department of Biochemistry, Central Research Institute of Medical Endiology, Leningrad.

(LIVER radiation eff.)

(GLYCOGEN metab.)

REYLINH, IT YO

69

PHASE I BOOK EXPLOITATION

807/5435

Kiselev, P. N., Professor, G. A. Gusterin, and A. I. Strashinin, Eds.

Voprosy radiobiologii. t. III: Sbornik trudov, posvyashchennyy 60-letiyu so dnya rozhdeniya Professora M. N. Pobedinskogo (Problems in Radiation Biology. v. 3: A Collection of Works Dedicated to the Sixtieth Birthday of Professor M[ikhail] M[ikolayevich] Pobedinskiy [Doctor of Medicine]) Leningrad.

Tsentr. n-issl. in-t med. radiologii M-va zdravookhrananiya SSSR, 1960.
422 p. 1,500 copies printed.

Tech. Ed.: P. S. Peleshuk.


PURPOSE: This collection of articles is intended for radiobiologists.


COVERAGE: The book contains 49 articles dealing with pathogenesis, prophylaxis, and therapy of radiation diseases. Individual articles describe investigations of the biological effects of radiation carried out by workers of the Central Scientific Research Institute for Medical Radiology of the Ministry of Public Health, USSR. [Tsentral'nyy nauchno-issledovatel'skiy institut meditsinskoy radiologii Ministerstva zdravookhraneniya SSSR] during 1958-59. The following

Card 1/10

i i		= 69	:
	Problems in Radiation Biology (Cont.) S07/5435		
	topics are covered: various aspects of primary effects of radiation; the course of some metabolic processes in animals subjected to ionizing radiated organisms; morphologic charges in radiation diseased and reparation and regeneration of tissues injured by irradiation. Some articles give attention to the effectiveness of experimental medical treation personalities are mentioned. References accompany almost all of the	ation; ase; tments.	
!	TABLE OF CONTENTS:		
	Foreword	3	:
	Gusterin, G. A., and A. I. Strashinin. Professor Nikhail Nikolayevich Pobedinskiy (Commemorating his Sixtieth Birthday)	5	
	Lebedinskiy, A. V. [Member, Academy of Medical Sciences USSR], N. I. Arlashchenko, and V. M. Mastryukova. On the Mechanism of Trophic Disturbances Due to Ionizing Rediation	n	į
† :	Zedgenidze, G. A., [Member, Academy of Medical Sciences USSR], Ye. A. Zherbin, K. V. Ivanov, and P. R. Vaynshteyn. Hormonal Activity of the Adrenal Cortex in Acute Radiation Sickness and the Effect of Desoxy-		
•	corticosterone Acetate on the Disease	17	
1	Card 2/10		
	the second secon		
	The state of the s		

		6
\$ \$	Problems in Radiation Biology (Cont.) SOV/5435	
	Manoylov, S. Ye., and B. A. Orlov. Use of the Spectroscopic Research Method in the Study of the Condition of Iron-Containing Compounds in Animal Organisms Irradiated With X-Rays	152
· }	Dumin, N. N., [Professor]. On Some Metabolic Disturbances in Lipides Due to External Whole-Pody Ionizing Irradiation	158
,	Keylina, R. Ya. Concerning the Problem of the Condition of Processes of Synthesis and Decemposition of Carbohydrates in the Animal Organism Subjected to Whole-Ekdy X-Ray Irradiation	165
† ! •	Shitova, Z. I. Changes in Respiration of the Skeletal Muscles Following Irradiatics by Radon of Different Segments of the Nervous System	173
	Mytareva, L. V. Effect of Ionizing Radiation on a Ferment of Glycolysis of Phosphohexcisomerase in Some Organs and Tissues of an Animal Organism Subjected to Whole-Body Irradiation	183
	Card 5/10	,
-		





KEYLIS-BORCK, V. I.

PA 233T84

USSR/Geophysics - Earthquakes

1950

"Determining the Dynamic Parameters of the Focus," V.I. Keylis-Borok

"Trudy Geofiz Inst" No 9 (136), pp 3-19

Considers a method of detg the source of an earthquake, approximately equiv to a focus, that is, attempts to develop a simpler approx method for selecting the type of source corresponding to the focus being studied.

233184

KEYLIS-BORCK, V. I.

PA 233T85

USSR/Geophysics - Earthquakes

1950

"Investigating Sources Equivalent to the Foci of Earthquakes," V. I. Keylis-Borok

"Trudy Geofiz Inst" No 9 (136), pp 20-42

Considers the relations between (a) certain peculiarities of the focus and (b) the functions which define the source approximately equiv to it. Studies the dipole with a moment as a source, which corresponds to a tectonic focus.

233185

"APPROVED FOR RELEASE: 09/17/2001

CIA-RDP86-00513R000721620005-8

A CONTRACT AND A SECURE AND A SECURE ASSESSMENT AS A SECURE OF A S KEYLIS-LÜRUK, V. I. owidat wi 21 Feb 50 USSR/Geophysics - Seismism (Earthquakes) "Dynamic Characteristics of a Centrum According to Seismic Observations," V. I. Keylis-Borok, Geophys Inst, Acad Sci USSR "Dok Ak Nauk SSSR" Vol "XX, No 6, pp 995-998 Applies several works of Acad V. I. Smirnov and S. L. Sobolev's school on theory of elasticity to interpretation of seismic data. Study of centra by method proposed would require seven or eight seismic stations located not further than 150-180 km from the centrum. Submitted 19 Dec 49 by Acad O. Yu. Shmidt. 165T26

Kayllis=70	DAOK, V. I.		~ MH 0 ~			1807
			crust. Analyzes: particular, the Rayleigh seismic (deceased).	E B O ct	H.	> # # #
		* •	R/G tic tei	tions generated by boundary of paralle half space. Comput characteristics of		USSR/Geophysics "Superficial Was Balf Space," V. Acad Sci USSR
	:		sed a	cte gas ge	F	Geophysi erficial Space," Sci USSR
			hys nalt r, r, sei	ris.		oph ici
	:		vze, the	tic	,	Geophysics erficial Wa Space, " V. Sci USSR
	÷ .		7 E 8 1	E E E E E E E E E E E E E E E E E E E	Ser	
		-	some e. nature	or J	န္	H (s)
			8 6 6 E	pertie sinus el pla tes di tayer	. 8	in Key
			St in plot	len lan	୍ ନୃ	116
		•		s of oidal ne la spers	· of	Seismology in a Laye Keylis-Bo
			sics - Seismology (Contd) yzes some explt data on serm the nature of Rayleigh and p smic waves. Submitted by L.	pared by sinusoidal coperated by sinusoidal coperated plane layer Computes dispersion classer to	19	Seismology in a Layer Lying Keylis-Borok, Geo
			ntd gh s	superf concd concd yer ly lon and to av	No	r y
	•		d) set and by I	v t	'n	95 C
	. · · · ·		ophysics - Seismology (Contd) Analyzes some explt data on serimic lar, the nature of Rayleigh and pseu n seismic waves. Submitted by L. S. ed).	sinusoidal concd force lying on es dispersion and frequest dayer corr to av terres	đđ	on I
	4			ced by sinusoidal concd force on frearellel plane layer lying on elast Computes dispersion and frequency cs of layer corr to av terrestrial	Ak Nauk, Ser Geog 1 Geofiz" No 2, pp 17-39	- Seismology Mar/ es in a Layer Lying on Elastic I. Keylis-Borok, Geophys Inst,
-	س		Mar/Apı waves; ido- Leyben	las ncy ría	39	15 t
	180763		Apr	rated by sinusoidal concd force on free of parallel plane layer lying on elastice. Computes dispersion and frequency stics of layer corr to av terrestrial		Mar/Apr stic nst,
	නු	1	Mar/Apr 51 waves; in do- Leybenzon	lla- free stic y al		Z Z
	44 T. 4 T. 1			L	• :	H
		. F.,	January Company			

USSR/Geophysics - Earthquake Centers Nov/Dec 51	
"Graphic Methods for Calculating the Dynamic Parameters of Earthquake Seats," V. I. Keylis-Borok	
"Iz Ak Nauk SSSR, Ser Geofiz" No 6, pp 11-27	
Considers graphic methods for detg the parameters of a source approximately equiv to the site (focus) of an earthquake. Gives extensive formulas, graph, and tables for subject detn.	
199172	

KEYLIS-BOROK, V. I.

USSR/Geophysics - Epicenter Azimuth


1952

"Problem of Determining the Azimuth to the Epicenter," V. I. Keylis-Borok, S. D. Kogan

"Trudy Geofiz Inst, Ak Nauk SSSR" No 14 (141), pp 21-28

Describes a method of detg the azimuth to the epicenter according to observations of transverse waves.

230165



DONON, V.I.

KEYLIS-BOROK, V. I.

"Problems of Propagation of Stationary Oscillations in a Layer Located Between Two Semispaces," Tr. Geofiz. in-ta AN SSSR, No 20, 1953, pp 20-36

Dispersion and resonance of stationary axially symmetrical, interfering oscillations generated by a centered force in a flat parallel layer to roots of frequency equations. The dependence of the amplitude of layer is computed in a few examples.

RZh Fiz. No 3, 1955


MEYLIS-DONON V.I., KOGAN, S.D.

Methods of seismic observations in an epicentral zone and their interpretations. Trudy Geofis.inst. no.21:27-48 153. (MLRA 7:5) (Seismology-Observations)

----- DUKUK, V. I. CIA-RDP86-00513R000721620005-8"

APPROVED EQRand Results of Earthquake Mechanisms Investigations," one of the Combusian Page 1051.

Evaluation, B-86198 and 86204, 30 Jun 55



Investigation of focal stresses of the Khait epicentral zone.
Trudy Geof.inst. no.25:113-123 '54. (MLRA 7:12)

(Khait--Seismology)

REILIS-BOROK, V.I. Propagation of waves in multilayer semi-spaces. Dokl.AH 888R 95 no.4:733-735 Ap 154. (Wibration) (Seismometry) (Vibration) (Seismometry)

KAYLIS-BOROK, V.I.

"Methodology and Results of Researches on Mechanisms on Earthquakes."

SO: Soviet Academy of Science Proceedings, No.6, March Issue 1955; A-40687

KEYLIS-BOROK B.I., prof. and KOGAN, S.D., Candidate of Physicomathe-matical Sciences

"Investigation of the mechanism of earthquakes", a paper given at the 50th Anniversary Session of the Seismic Station "Pulkovo", 25-29 Sep 1956, Leningrad.


SUM. 1322

KEYLIS-BOROK, V.I.

Relation of point and area sources. Izv.AN SSSR.Ser.geofiz. no.4: 404-409 Ap '56. (MLRA 9:8)

1. Akademiya nauk SSSR, Geofizicheskiy institut. (Seismology)

APPROVED FOR RELEASE: 09/17/2001 CIA-RDP86-00513R000721620005-8"



	SACTIONS OF THE DESCRIPTION OF THE PROPERTY OF	A CONTRACTOR OF THE PROPERTY O
KAWAS ROPIN 1	/ T	
KEYLIS-BOROK, I	. Bararua I sa	
	· BALAKINA, L.M.	
	X(10) PRANE I BOXX EXPLOIPMENT BOX7.6	69
	Abademiya menk 888R, Komitet po geodemii i geofinike.	
	Texisy foliadov as XI General'may assembleys Heabhmarodnego geodesiches i geofisicheskogo sequesa. Heabhmarodneye assotsiatsiya sequenclogii fiziki medr semii (Abstracts of Reports Submitted to the II General International Jaious of Geodesy and Geophyses. The International Locations of Sciencing and Physics of the Earth's Interior) Moscow, 1997. 108 ps. /Farallel texts in Emsim and Engli	
	No additional contributors mentioned	菱
	FREPOSE: This booklet is intended for geographicists, especially those iming in scientology.	1.4
	COVERAGE: This collection of articles deals with the structure and comp of the Earth and phenomena related thereto. The majority of the arti- serm studies of surthqualus and, seismic invest, Other articles cover ture of the Earth's crust and mountain roots; the classic properties at high pressures; the piescollectic affects of the classic properties.	cles con-
	at high preserve; the piesoclastric effect of rocks and the method a modelling in testmophysics. The collectim size contains ratiole on the strike thermi history, the ninvestenic method of treeing others.	
	Volarovich, N.P. and R.I. Furthemento. Pisonoelsetric Riffeet of & he	29
	Veytman, P.S., I. P. Kominskeys, and Ts. V. Risnichenko. Her Bridene on the Structure of the Earth's Cruet and Noustain Scote in Central Asia From Seisnie Dopth Sounding Data	
	Gaovekly, M.V. Nethod of Modelling in Testonophysics	N A
	Gorahkov, G.P. Science Intensity Regions of Asia	π }
	Derrier, B.I. Pronted Security Degices of Asia	19
	Derrier, B.I. Paysical Properties of Solid Bodies at Righ Processes	u
	Esplis-Borok, Val. Investigation of Marthquake Hechanism	₩ 🖟
	Reylis-Brok, V.I. Bysamic Sethods of Investigating the Sorth's Crust Fraction, Sets) Fraction, Potes) Reylis-Brok, V.I. Bysamic Sethods of Investigating the Sorth's Crust Fraction, Sets)	1
	Exrus, To.T. Absorption of Elastic Pures in Rollin	s .
	Care 1/9	55
	หลังของเกิดสำหรับแล้วสามารถ เกิดสาร์ แล้ว รับเกิดสาร์ และสาร์ และสาร์	Contract Co
	· 中国等人的 唯《阿特·SP·斯··································	

GOTSADZE, O.D.; KIRILLOVA, I.V.; KOGAN, S.D.; KUKHTIKOVA, T.I.;

MALINOVSKAYA, L.N.; SORSKIY, A.A.; KEYLIS-BOROK, V.I.,
doktor fiziko-matematicheskikh nauk, otvetstvennyy redaktor;
ZAYTSEV, L.P., redaktor izdatel'stva; EZ. V.V., redaktor
izdatel'stva; SHEVCHENKO, G.N., tekhnicheskiy redaktor.

[Investigation of the mechanism of earthquakes] Issledovanie
mekhanizma zemletriasenii. Moskva, Izd-vo Akademii nauk SSSR,
1957. 148 p. (Akademiia nauk SSSR. Geofizicheskii institut.
Trudy, no.40).

(Seismolegy)

KEYLLS-BOROK, V. I., REZNICHENKO, Yu. V., BELOUSOV, V. V. MAGNITSKIY, V. A., and LYUBIMOVA, Ye. A.,

"Seismological Problems and Questions Concerning the Physical Structure of the Earth's Deposits."

paper presented at the XIth General Assembly of the Int'l. Union of Geodesy and Geophysics, Toronto, Canada, 3-14 Sept. 1957, (Izv. Ak Nauk SSSR - Ser. Geog. 1958, No. 2, pp 3-8 [USSR]).

APPROVED FOR RELEASE: 09/17/2001 CIA-RDP86-00513R000721620005-8"

The Harmon Commence of the Com

KEYLIS-BOROK, V. I.

"On a Method He Developed for Determining the Elements of the Occurrence of Recent Faults Found in the Depths and Causing Earthquakes, and an Explanation of the Direction of the Displacement of their Walls," paper presented at the First All-Union Conference on Tectonphysics, Moscow, 29 January through 5 February 1957.

Institute of Physics of the Earth, Academy of Sciences USSR

Sum 1563

KEYLIS-BOROK, V.I.; MALINOVSKAYA, L.N.

Dislocations in the foci of weak earthquakes in northern Tien Shan.

Biul. Sov. po seism. no.3:118-122 '57.

(Tien Shan--Earthquakes)

KEYLN-BOROK

AUTHOR: Keylis-Borok, V. I. 49-4-3/23

TITLE:

On the theory of waves generated during shear. (O teorii voln, voznikayushchikh pri sdvige).

PERIODICAL: Izvestiya Akademii Nauk, Seriya Geofizicheskaya,

1957, No.4, pp. 440-448 (ÚSSR)

用自作的 數4. 海國自然發展的基準

ABSTRACT: The relation is investigated between a point source of elastic waves (a model simulating the focus of an earthquake) and the static displacements which occur after passage of these waves. An analysis is made of the static field, described by means of the dislocation theory as formulated by Love for shear dislocations, and the possibility of application of this theory to simulating such of focion on models. In the case of correct analysis, the conclusions derived from the dislocation theory are in agreement with generally accepted conceptions relating to a shear wave in the shape of a dipole with a moment. Vvedenskaya, A.V. (Ref.4) attempted to find a shear model by means of the Love dislocation theory, considering as a model of the elementary shear two perpendicular dipoles with a moment as indicated in Fig. 2 and not one dipole with a Card 1/5 moment. The dependence of the intensity of the source

ARPROVED FOR RELEASES Q9/17/2901duriGIA-RDP86-0051/3R40972/1620005-8"

on the time is thereby so chosen that the main part of the dislocations is represented by the Dirac 5-function, referred to in this paper as the Nabarro source; the displacements described by the Nabarro formula (Ref.5) are integrated along a circle which simulates the sliding plane in the tremor and it is assumed that the obtained expressions describe waves which occur during shear, taking into consideration the presence of fault. It is shown in this paper that Nabarro did not generalise the theory of dislocations for the dynamic case and introduced certain assumptions without any proof; extension by Vvedenskaya of the respective interpretation led to erroneous results, even if it is assumed that the assumptions of Nabarro are correct. Application of the theory of dislocations in modelling of earthquakes is fundamentally based on the assumption that residual deformations are absent in the focus; even for this assumption, the results obtained from the theory of dislocations are practically equal with those obtained in earlier work of the author (Refs.1 and 2). The theory of dislocations in the sense expressed by Love describes the state of a continuous

Card 2/5 medium and permits determination of a source with static forces with equivalent dislocations, i.e. sources which

KEYLIS-BOHOK, V.I.; KOGAN, S.D.

Investigating the mechanism of earthquakes. Biul. Sov. po seism. no.6:96-99 '57. (MIRA 11:3)

1. Institut fiziki Zemli Akademii nauk SSSR, Moskva. (Earthquakes)

计写话 计概号 医乳

SOV/169-59-3-2207

Translation from: Referativnyy zhurnal, Geofizika, 1959, Nr 3, p 21 (USSR)

AUTHORS: Keylis-Borok, V.I., Kogan, S.D.

TITLE:

TARPES P

An Investigation of the Earthquake Mechanisms

PERIODICAL: Byull. Soveta po seysmologii AS USSR, 1957, Vol 16, pp 96-99

ABSTRACT: This is a brief survey of the work of Soviet authors on the mechanisms of earthquakes. The analysis of the investigation results of about 300 earthquakes in the main seismoactive zones

of the USSR and adjacent regions leads to the following preliminary conclusions: 1) The predominant properties of dislocations in the earthquake source vary sharply with the transition from one tectonic section to another one. There is also no direct link of dislocations in individual sources with local tectonic surface fractures. 2) In regard to the strike of the basic structures, the predominant strike of dislocation is transverse for the My part of the Recific Ocean December the Albertales.

for the NW part of the Pacific Ocean, Dagestan, the Akhalkalaks-koye highlands, and longitudinal for the Ashkhabadskiy rayon.

Card 1/2 But in the majority of zones (Garmskaya oblast', Tien Shan,

An Investigation of the Earthquake Mechanisms SOV/169-59-3-2207

Shemakha), two predominat strikes are distinguished, the longitudinal and the transverse strikes. A gradual change of the strike of dislocations in the foci from transverse to longitudinal is to be noticed in the Prikazbekskiy rayon of the Caucasus with increasing distance from the zones of transverse fractures. 3) The horizontal components of the displacements in the foci have great intensity. 4) The vertical components of the motion in the foci of some districts (West-Turkmenistan, Caucasus, Tien Shan) have a peculiarity consisting in a lift of the reverse wing, located on the side of the depression. In general, the results adduced coincide with the neotectonic data for a number of regions, where recently formed surface dislocations differ from the basic, older structures, approximately in the same manner as the dislocations in the earthquake foci described by the authors.

S.D. Kogan

V

Card 2/2

KEYLIS-BOROK, V. I.

"The relation of the Area of Disruption to the Energy of Seismic Waves."

report presented at a meeting of the European Commission of the International Association of Seismologists and Earth Physicists, Utrecht, Holland, 8-12 April 1958. (Izv. Akad. Neuk SSSR, Ser Geofizicheskaya, 1958, No. 11, pp 1418.)

Ceffin - BOROK, U.Z.

24(6)

PHASE I BOOK EXPLOITATION SOV/2250

Akademiya nauk SSSR. Institut fiziki zemli

Nekotoryye voprosy mekhaniki deformiruyemykh sred (Some Problems in the Mechanics of Deformable Media) Moscow, Izd-vo AN SSSR, 1959. 219 p. (Series: Its: Trudy, Nr. 2 /169/) Errata slip inserted. 2,000 copies printed.

Ed.: V.A. Magnitskiy, Doctor of Technical Sciences; Ed. of Publishing House: V.A. Kalinin; Tech. Ed.: Yu. V. Rylina.

PURPOSE: This book is intended for engineers and geophysicists concerned with problems of deformations.

COVERAGE: This collection consists of eight articles on the mechanics of deformations in solid plastic media as applied to the solution of geophysical and engineering problems. No personalities are mentioned. References appear at the end of each article.

TABLE OF CONTENTS:

Card 1/5

APPROVED FOR RECEASE) 09/17/2001 CIA-RDP86-00513R0003731620005-8"

Gurevich, G.I., and A.L. Rabinovich. Relation Between Stresses and Displacements in Large Deformations for the Case of a One-dimensional Problem

In the analytical study of geometrical and kinetic deformations of elastic and residual nature, which are of signficance in attenuation and dispersion of seismic waves, the authors derive general equations of motion.

Gurevich, G.I. Relation Between Stresses and Displacements in Large Deformations for the General Case of a Three-dimensional Load

The author considers the application of Maxwell's equation to a case of a residually deformed solid-liquid body which can be considered as a "massive" one and to which the usual formulas of the theory of elasticity are applicable.

Gurevich, G.I. Generalized Maxwell Equation for Three Measurements Taking Into Consideration Small Elastic Aftereffect Deformations 60 In the study of rock behavior in cases of static and dynamic

Card 2/5

27

12

3

Some Problems (Cont.)

SOV/2250

Petrachen', K.I. Ogurtsov.

三型产品建筑设置建筑设置建设。在外位

Khaykovich, I.M. Beam Method of Computing the Wave Intensity in a Relaxing Medium With a Large Relaxation Time 179

The author refers to various scientists offering the solution of nonstationary problems in the theory of elasticity leading to the determination of the intensity and the force of reflected waves. He introduces a so-called beam method for computing the propagation of a wave in nonideal elastic media. The following names are mentioned: G.I. Petrashen', V.M. Babich, G.O. Gurevich.

Sherman, D.I. Problem of the Stressed Condition of a Semiplane Without External Load and With Two Sunken Circular Orifices 187 The article discusses the distribution of stresses caused by gravity in media weakened by holes or openings. The problem is of interest in analyzing the rock pressure in the neighborhood of shaft openings and for the study of seismic conditions.

Card 4/5

SOV/49-59-1-4/23

AUTHORS: Keylis-Borok, V. I. and Stesin, I. M.

Dispersion of the Raleigh Waves in a Two-Layer Model TITLE: of the Earth's Crust (Dispersiya releyevskikh voln

v dvukhsloynoy modeli Zemnoy kory)

PERIODICAL: Izvestiya Akademii Nauk SSSR, Seriya Geofizicheskaya, 1959, Nr 1, pp 27-31 (USSR)

ABSTRACT: In order to improve the data for the Earth's crust calculated by the dispersion method, a two-layer dispersion method is proposed. The calculation is based on the half-space z 0 covered by two layers:

 $0 > z > -m_1$ and $-m_1 > z > -(m_1 + m_2)$.

The base and the two layers are composed of a homogeneous medium which is also isotropic and ideally elastic. The phase velocity v of the Rayleigh waves in such a medium can be represented by Eq.(1), where mo - thickness of q-layer,

- velocities of longitudinal and transverse

waves respectively,
- a constant (q = 1, 2, 3 denote top, middle layers
and base respectively).

SOV/49-59-1-4/23 Dispersion of the Radeigh Waves in a Two-Layer Model of the Earth's Crust

The relationship of these values is calculated from the

$$\sigma_{\rm q} = \mu_{\rm q+1}/\mu_{\rm q}; \ \alpha_{\rm q} = \sqrt{1 - {\rm v}^2/a_{\rm q}^2}; \ \beta_{\rm q} = \sqrt{1 - {\rm v}^2/b_{\rm q}^2}; \ \gamma = 2 - {\rm v}^2/b_{\rm q}^2;$$

$$s = 2(\sigma_1 - 1) + v^2/b_1^2$$
, $c = 2(\sigma_1 - 1) - \sigma_1 v^2/b_1^2$; $d = c + v^2/b_1^2$;

$$X_{i_{\alpha}} = \operatorname{ch} \alpha_{i} \xi_{m_{i}}; Y_{i_{\alpha}} = \operatorname{sh} \alpha_{i} \xi_{m_{i}}; X_{i_{\beta}} = \operatorname{ch} \beta_{i} \xi_{m_{i}}; Y_{i_{\beta}} = \operatorname{ch} \beta_{i} \xi_{m_{i}}.$$

(ξ - wave characteristic, ξ = p/v, p - rotational frequency). The dispersion surface is equal to $v/b_3 = f(\lambda/m_1, \lambda/m_2)$, where $\lambda = 2\pi (b_3/p)$ - the length of the transverse wave with frequency p. It is divided into successive parts according to the different harmonics (Refs 2,4,5). For computation on hand machines Card 2/4 it is better to find the solution of 5 m1 with given

Dispersion of the Radeigh Waves in a Two-Layer Model of the

 v/b_3 and f, m_2 . The results obtained by means of the electronic machine BESM, Ac.Sc., USSR are shown in Figs. 1-4. The constants used were those for the Central Asia (Refs 6,7) $a_3/b_3 = 1.85$, $a_2/b_3 = 1.46$; $a_1/b_3 = 1.28$, $b_2/b_3 = 0.88$, $b_1/b_3 = 0.76$, $\sigma_1 = 1.48$, $\sigma_2 = 1.48$ (a_3 , b_3 - the outer surface). Fig.1 shows the data of the dispersion surface: the curves of the phase v and the group velocity v of the first harmonic for various values of v are arc tg v and v and the group velocity v of the first harmonic for various values of v are arc tg v and v and v and v are 0.9265 for v and v are 0.9265 for v are 0.6923 for v are 0.8011 for v and 0.7 for v are 0.8011 for v and 0.8011 for v and 0.8011 for v are 0.8011 for v are 0.8011 for v and 0.8011 for v are 0.8011 for v are 0.8011 for v and 0.8011 for v are 0.8011 for v and 0.8011 for v are 0.8011 for v and 0.8011 for v are 0.8011 for v are 0.8011 for v and 0.8011 for v are 0.8011 for v are 0.8011 for v are 0.8011 for v and 0.8011 for v are 0.8011 for v are 0.8011 for v are 0.8011 for v are 0.80

Card 3/4 group velocity has a second minimum in high frequencies (Ref 1). It can be seen from Fig.1 that when the value

国际的基础的

SOV/49-59-1-4/23


Dispersion of the Radeigh Waves in a Two-Layer Model of the Earth's Crust

of m_1/m_2 increases, the equivalent value $(m_1 + m_2)$ decreases. This is explained by a decrease of the wavelength due to $b_1 < b_2$. In order to determine m₁ and m₂ by means of the curves (Fig.1), it is necessary to find the dispersion V in the homogeneous region so that the respective value of b₃ can be established. (The notations of Fig.1 are as follows: 1 - phase and 2 - group velocities, 3-6 - experimental data for Raneigh waves in various localities (Ref 1)). There are 4 figures, 1 table and 8 references, 5 of which are Soviet, 3 English.

ASSOCIATION: Akademiya Nauk SSSR Institut fiziki Zemli

(Ac.Sc. USSR, Institute of Earth Physics)
SUBMITTED: September 10, 1957

Card 4/4



66303

3.9100

SOV/49-59-11-1/28

AUTHORS:

Keylis-Borok, V. I., and Monin, A. S.

TITLE:

Magneto-elastic Waves and the Boundary of Earth Core

PERIODICAL: Izvestiya Akademii nauk SSSR, Seriya geofizicheskaya,

1959, Nr 11, pp 1529-1541 (USSR)

ABSTRACT: The authors describe results of their investigations of the dispersion, damping, polarization and origin of magneto-elastic waves. The damping of waves with an increase of the field strength Ho attains a certain maximum value from which it gradually tends to 0. Two types of waves can occur in a strong field: slow waves with a velocity between the transverse and longitudinal elastic waves, and fast waves with a velocity proportional to $H_{\mathbf{O}}$. An intense mechanical vibration can only be caused by slow waves. The characteristics of the latter, being similar to longitudinal or transverse waves, depend on the direction of their propagation and are not affected by the initial pulse. The observed range of velocities of seismic waves in a stratum D" can be explained by a linear increase of the gradient k/P and by a decrease of u/P (see table) due to compensation by the magnetic field. The existence of transverse waves in the core can be Card 1/2 explained by a presence of the magnetic field

APPROVED FOR RELEASE: 09/17/2001

CIA-RDP86-00513R000721620005-8" SUV/49-59-11-1/28

Magneto-elastic Waves and the Boundary of Earth Core

unless there is a layer 20 to 80 km thick at its surface with an increased resistance (to \sim 0.4 x 10⁻⁸ sec) or field with a strength $\sim 10^9$ to 10^{10} there is a radial oersted. An increase of the phase velocity v of longitudinal and transverse (dashed lines) waves is illustrated in Fig 1, where the numbers on curves denote a squared strength of the field ϕ . Fig 2 shows the polarization of vibration in a weak field. Fig 3 illustrates the phase velocity of vibrations in a strong field. Fig 4 gives the coefficients of damping with distance in a weak field where the magnitudes of $e^{-K_1} \times \omega/a$, $e^{-K_1} \times \omega/a$ determine damping with a distance x of longitudinal and transverse waves respectively. Thanks are conveyed to G. S. Golitsyn, V. A. Kalinin and R. Khayd for their assistance. There are 4 figures, 1 table and 12 references, 6 of which are Soviet and 6 English.

ASSOCIATION: Akademiya nauk SSSR, Institut fiziki Zemli (Academy of Sciences USSR, Institute of Physics of Earth)

SUBMITTED: March 26, 1959

3\S brev

PHASE I BOOK EXPLOITATION

SOV/4521

POSTAL PARTIE PROPERTY OF THE PROPERTY OF THE

Keylis-Borok, Vladimir Isaakovich

Interferentsionnyye poverkhnostnyye volny (Interference Surface Waves) Moscow, Izd-vo AN SSSR, 1960. 194 p. Errata slip inserted. 3,000 copies printed.

Sponsoring Agency: Akademiya nauk SSSR. Institut fiziki zemli im. O. Yu. Shmidta.

Resp. Ed.: D.I. Sherman; Ed. of Publishing House: T.B. Yanovskaya; Tech. Ed.: T.V. Polyakova.

PURPOSE: This book is intended for geophysicists, seismologists, meteorologists, and oceanographers.

COVERAGE: This is a study of oscillations of a layered elastic medium (plane-parallel, uniform, isotropic, and ideally elastic layers) from a single viewpoint. The general physical and mathematical description of the oscillations is suitable for direct analysis of the mathematical derivation and for studying the solutions together with the system of houndary conditions, without resorting to cumbersome solutions of elastic theory problems. This approach made it possible to obtain general results for any number of layers and for practically any nonsymmetric,

Carrie / 0

Card 0/8

GAMBURTSEV, Grigoriy Aleksandrovich, akademik [decessed]; RIZNICHENKO, Yu.V., red.; MOLODEMSKIY, M.S., red.; BERZON, I.S., doktor fiz.-mat.nauk, red.; KEYLIS-BOROK, V.I., doktor fiz.-mat.nauk, red.; LYAPUNOV, A.A., doktor fiz.-mat.nauk, red.; YEPINAT'YEVA, A.M., kand.tekh.nauk, red.; KOSMINSKAYA, I.P., kand.fiz.-mat.nauk, red.; STARODUBROVSKAYA. S.P., mladshiy nauchnyy sotrudnik, red.; BERKGAUT, V.G., red.izd-va; MARKOVICH, S.G., tekhn.red.

[Selected studies] Izbrannye trudy. Moskva, Izd-vo Akad.nauk SSSR, 1960. 461 p. (MIRA 13:7)

1. Chleny-korrespondenty AN SSSR (for Riznichenko, Molodenskiy). (Prospecting-Geophysical methods)

APPROVED FOR RELEASE: 09/17/2001 CIA-RDP86-00513R000721620005-8"

the literature, indicate that without preliminary analysis of a contact of observations there is no reason for overall study is a new region of the mechanics of earthquakes, or the mass processing of any system of observations for my purpose whatsowers. The principle at the bare of 'no proposed method for studying the prevalent attributes and duy, shifts of discontimities, stragen be applied in the solution of a number of other problems. No personalities are mentioned.	Til Darrich, F.L., and O.L. Barlora. Generalization of Data on the carried to Extraplace to the characteristic of observations to any creatized parameters of the schools for perhaps of the schools of extraplace is in any carried and parameters of the schools of extraplace is in a carried was schools appeared to be extraplaced in the carried as a carried and carried as each for the fall those Baltham countain reage (Tarcacratic). The color conclude that the method described is comberness, but that it is next (field by the remains of each idea should necessary be a that it is next (field by the remains of each idea of this study, based on observation of the fall thinks and the color of the study, based on observation of the fall thinks and on the color of the study as the classification of the fall of the color of the study as the classification of the fall thinks and on the color of the study of easiers. And on the color of the fall of the color of the study of the color of the study of easiers.	comy tenders so with from law by I.D. Ebone little, "The problem when craylically. The server in the calculation of equitalists duringly is 10% of the maximum value. The errors in great theretic much make the control of equitalists. Several theories as to the monutions mass distribution when a service the maximum control of the contro	e plice	agmitally, Doctor of Te- ch. Ed.: E.G. Thbond lection of articles is ta. The properties of the Properties and tentals articles of victors in the Table of collection have been a	PRACE I BOOK EXTRODISHING SOY/ASO Absoluty near SECE. Institut finite Semil Topping teorestichenkay seymologit i fisht semiph medr (Problems in the Theory of Settemblogy and Restes of the Kuthin Interior) boscov, 1960, 177 p. (Series: Its: Trudy, no. 11 (178)) Erreta slip inserted, 1,700 copies print- pomoring Agency: Alademiya near SECE, Institut fishat Esmil incol 0, To.	
---	---	--	---------	---	--	--

S/169/61/000/011/006/065 D228/D304

AUTHOR:

Keylis-Borok, V.I.

TITLE:

Studies of the dislocations at earthquake foci

PERIODICAL:

Referativnyy zhurnal. Geofizika, no. 11. 1961 14.15. abstract 11A141 (V sb. Probl. tektonofiziki, M., Gos.

geoltekhizdat, 1960, 309)

TEXT: A method permitting determination of the properties of dislocations at an earthquake focus from seismic data (the mechanical type of fracture, the dip and strike of the plane of fracture, the direction of movement) was elaborated. About 300 earthquakes were investigated in the main seismically-active zones of the Soviet Union and adjoining regions: the north-west edge of the Pacific Ocean, the Hindukush, the Pamirs and Tyan Shan Kopet-Dag and Western Turkmeniya, and the Caucasus. On the whole, the dislocations at the foci have approximately the same similarities to and differences from the main tectonic structures as those which have been formed in recent times and which are distinguishable in neo-

Card 1/2

Studies of the dislocations at ...

S/169/61/000/011/006/065 D228/D304

tectonics. Dislocations with a transverse strike and with intense horizontal components of movement are evidently noted more frequently at the foci than is the case in surface structures. Vertical movements at the foci of a number of areas are characterized by the uplift of the limb of the fracture which has been turned to the side of the tectonic depression. This agrees with the data of nestectonics and with some results of abyssal seismic sounding. [Abstractor's note; Complete translation].

Card 2/2

S/169/61/000/010/009/053 D228/D304

AUTHORS:

Bune, V. I., Gzovskiy, M. V., Zapol'skiy, K. K., <u>Keylis-Borok</u>, V. I., Krestnikov, V. N., Malinovskaya, L. N., Nersesov, I. L., Pavlova, G. I., Rautian, T. G., Reysner, G. I., Riznichenko, Yu. V., and Khalturin, V. I.

TITLE:

Methods of the detailed study of seismicity

PERIODICAL:

Referativnyy zhurnal, Geofizika, no. 10, 1961, 12-13, abstract 10A144 (Tr. In-ta fiz. Zemli AN SSSR, no. 9, 1960, 327 p.)

TEXT: The Tadzhik complex seismologic expedition was organized with the aim of studying the nature of earthquakes and the conditions of their genesis. The most seismically-active zones of the USSR (Garmo and Stalinabad) were chosen as the work areas. The specific conditions of working and processing the data demanded the development of special systems of observation and methods of interpretation. The large amount of recorded

Card 1/6

S/169/61/000/ 010/009/053 D228/D304

Methods of the detailed ...

seismic phenomena permitted the use of statistical methods for studying their distribution in space and time; these methods, in their turn, provided the basis for introducing the quantitative indices of the seismicity characteristics of the seismically-active areas. The actual seismic observations were closely coordinated with geologic investigations, and this provided the possibility of exposing the tectonic basis of the seismic phenomena. A general review of the work area is given in Chapter 1, and concise data on major earthquakes are cited together with the general position of the expedition stations. A description of the standard main and auxiliary apparatus used at the stations, and also the layout and description of newly developed equipment -- including an automatic seismic station with a magnetic memory -- is cited in Chapter 2. The methods developed and utilized in the expedition for studying the crust's structure in the area under investigation from the records of nearby earthquakes are described in Chapter 3. Horizontal and vertical hodographs were constructed. resulting material enabled the crust to be represented as a one-layer mass

Card 2/6

Methods of the detailed ...

5/169/61/000<mark>/010/009/053</mark> D228/D304

with a longitudinal-wave velocity of 6.0 - 6.1 km/sec. At the Mohorovicic boundary, the velocity suddenly changes to 8.0 km/sec. and then somewhat decreases, but at a depth of 300 km it subsequently increases to 9.2 km/sec. These data underlay the construction of isochrone charts used to localize the epicenters and to determine the focal depths. The isochrone charts were constructed with an account of the heterogeneity of the work area's geologic structure and the peculiarity of the seismic stations' location. This enabled the precision of hypocenter localization to be substantially increased, reducing it to 1 - 2 km at the center of the work area's topographic map. In Chapter 4, the definition of the concept of seismic energy at the focus is given, and the basic formulas are derived for its calculation. On the basis of experimentally obtained laws for the dying out of energy with distance, nomographs were constructed to determine practically the energy at the focus from the records of nearby earthquakes. Appraisal of the precision of calculation of the energy in relation to different factors shows that it may be determined accurately to the order of its magnitude. In this connection, the value K = lg E j.

Card 3/6

APPROVED FOR RELEASE: 09/17/2001 CIA-RDP86-00513R000721620005-8"

S/169/61/000/010/009/053 D228/D304

Methods of the detailed ...

is introduced for characterizing the energy class of earthquakes. The value of K is compared with the earthquake magnitude M. The study of the iso-energy lines shows that the different degrees of the dying out of seismic energy along and across the strike of geologic structures exert a decisive influence on the form of the isoseisms. In Chapter 5, the frequencies of seismic vibrations are studied -- in relation to the earthquake energy, the distance from the source, the geologic conditions at the point of observation and at the hypocenter, etc .-- from recordings at both the customary stations and a special 4NCC (ChISS) seismic-station intended for frequency analysis of seismic waves directly at their place of registration. A detailed description is given for the frequencyselective seismic-station 4%(C-1954 (ChISS-1954) and for the results of the investigation of its recordings. Certain epicentral zones with an anomalous frequency are thereby revealed. The procedure for theoretically calculating the focal characteristics, and also for appraising these latter from empirical data, is given in Chapter 6. Several formulas are

card 4/6

Methods of the detailed ...

S/169/61/000/010/009/053 D228/D304

cited for determining the size of a focus in relation to its energy on the basis of different physical propositions. The dynamic parameters of the foci are determined; there appear to be definite predominant directions for both the strike and dip of the fracture planes. The characteristics of the seismic conditions of the Garmo and Stalinabad seismically-active regions -- both as a whole and in individual areas -- are quoted together with the variations in the parameters of the conditions in time. The quantitative expression of the seismicity during constant seismic conditions is determined by the seismic activity. The possibility is shown of constructing graphs of the recurrence of earthquakes from short observations of weak shocks, and methods are given for determining the period required to obtain the parameters of he seismic conditions with a pre-set precision in relation to the energy of the recorded earthquakes. The statistical constancy of the seismic conditions is determined by the so-called measure of dispersion of the frequency of earthquakes. A brief description of the area's stratigraphy and the history of its geologic development is given in Chapter 8. The structural schemes and descriptions of the most important


Card 5/6

S/169/61/000/010/009/053 D228/D304

Methods of the detailed ...

deep faults are cited. The contemporary structure of the Garmo area is depicted as two main regions: the alpine geosynclinal zone in the south and the activated epi-Hercynian platform in the north. In section, it is drawn as several steps of Paleozoic basement adjoining each other along deep faults. A comparison of the seismicity with the tectonics of the study areas is made in Chapter 9. The construction of maps of isolines of seismic activity and gradients of the rate of tectonic movements is recommended for appraising the connection between the seismicity and the tectonics. Methods are cited for constructing such maps. The congruence between these magnitudes is established for the regions under investigation, and areas with the maximum gradient values correspond to those with the highest values of seismic activity. 272 references. Abstracter's note: Complete translation.

Card 6/6



3,9300(1406,/019,1109,1327)
AUTHOR: Keylis-Borok, V. I.

32023 \$/619/60/000/015/003/004 D039/112

.

TITLE:

The difference of the surface wave spectra of earthquakes and underground explosions

SOURCE: Akademiya nauk SSSR. Institut fiziki zemli. Trudy, no. 15 (182), Moscow, 1960. Seysmicheskiy effekt podzemnykh vzryvov, 68-101

TEXT: The dependence of the surface wave spectrum of the Rayleigh and Love type on the spatial properties of the source is investigated for the case of the most simple explosion and earthquake models. The effect of the dimensions and shape of the source on the surface wave spectrum and the intensity distribution in this source in the case of a given frequency spectrum of perturbance stresses is studied. Based on formula

 $U \approx R_{A} \left(\frac{p}{v}, \phi\right) P_{U} \left(\frac{p}{v}, z_{o}\right) Q(p) \frac{1}{\sqrt{r}} e^{-i\frac{p}{v}r}, \qquad (2)$

where $R_{\rm A}$ is the spectrum of the source according to distance, and on other calculations in the article, the author compares the surface wave spectra Card 1/3

等的可以發展的

32023 S/619/60/000/015/003/004 D039/D112

The difference of the surface wave spectra ...

for the accepted earthquake and explosion models. Formula (2) shows that the difference in the surface wave spectra will be defined by the functions of R which are independent from the structure of the medium. The calculations show that the difference between the sources is mainly determined by their dimensions. The greater the dimensions, the longer will be the predominant surface wave period. At the same time, the dimensions of the earthquake focus are much larger than those of the explosion zone. Consequently, the surface wave periods during earthquakes are also much longer than during explosions. It was established that there is a great difference among the surface wave spectra, but no difference in the spectra of volumetric waves. During earthquakes the stress concentration is considerably smaller than in explosions of the same energy level. So, in explosions, the non-elastic deformation zone will be much smaller. The depth of the source of earthquakes is larger than that of explosions. The theoretical conclusions of the author agree with experimental data taken from the paper of S. D. Kogan, I. P. Pasechnik, and D. D. Sultanov (Ref. 7: "Dokl. AN SSSR", 29, No 6, 1959.). For an explosion of 1 kg of explosives (according to a report by D. A. Kharin), the seismic wave energy was 4 . 1017 erg and the destruction zone diameter ranged between 150 and 200 m. It was

Card 2/3

KEYLIS-BOROK, Vladimir Isaakovich; NERSESOV, Igor' Leonovich; YAGLOM, Akiva Moiseyevich; SADOVSKIY, M.A., otv. red.; MEDER, V.M., red. izd-va; YEPIFANOVA, L., tekhh, red.

[Methods of evaluating the economic effect of earthquakeproof construction]Metodika otsenki ekonomicheskogo effekta seismostoikogo stroitel'stva. Moskva, Izd-vo Akad. nauk SSSR, 1962.
45 p. (MIRA 16:3)

1. Chlen-korrespondent Akademii nauk SSSR (for Sadovskiy). (Earthquakes and building)

BARLAS, V.Ya.[translator]; KEYLIS-BOROK, V.I., red.; RIZNICHENKO, Yu.V., red.; PANTIYEVA, V., red.; DOTSENKO, V., tekhn. red.

[Underground nuclear explosions]Podzemnye iadernye vzryvy.
Moskva, Izd-vo inostr. lit-ry, 1962. 247 p. (MIRA 15:8)
Translated from the English.
(Underground nuclear explosions)

APPROVED FOR RELEASE: 09/17/2001 CIA-RDP86-00513R000721620005-8"

KARUS, Ye.V., kand. fiz.-mat. nauk, red.; KEYLIS-BOROK, V.I., doktor fiz.-mat. nauk, red.; IONEL', A.G., ved. red.; YAKOVLEVA, Z.I., tekhn. red.

[Problems in seismic prospecting]Problemy seismicheskoi razvedki; sbornik statei. Moskva, Gostoptekhizdat, 1962. 225 p. Translated from the English and French. (MIRA 15:12) (Seismic prospecting)

APPROVED FOR RELEASE: 09/17/2001 CIA-RDP86-00513R000721620005-8"

42053

3.4300

S/619/62/000/020/001/001 1046/1246

AUNHORG:

Roylin-Borck, V.1. and Malinovskaya, L.N.

Tirli:

Identification of explosions from the arrival of body waves

Jeuros:

Akademiya nauk C.C.R. Institut fiziki Zemli. Trudy, no. 20(187), 1962. Voprosy teoreticheskoy seysmologii i fiziki zemnykh nedr, 117-124

TEXT: In earthquakes, both rerefection and compression waves are redicted from the focus, whereas in explosions only compression waves are generated. There is, however, a small but finite probability (of the order of 1%) that even in case of earthquakes only compression waves are recorded at all observation points. In this case, earthquakes are distinguished from explosions by the sigm of the first arrivals of the transverse SV waves as recorded on stations within a sufficiently wide range of azimuths: negative SV arrivals are characteristic of earthquakes, whereas positive SV arrivals characterize an explosion. There are 8 figures.

Card 1/1

KEYLIS-HOROK, V.I.; YANOVSKAYA, T.B.

Relation between the spectra of surface waves and the depth of the source in the earth's core. Izv. AN SSSR. Ser. geofiz. no.11:1532-1539 N '62. (MIRA 15:11)

1. Institut fiziki Zemli AN SSSR.
(Seismic waves—Spectra)

Report submitted to the Symposium on Results of the IGY-IGC (Intl. Geophysical Year), Los Angeles, California 12-16 Aug 1963.

PYATETSKIY-SHAPIRO, I.I.; ZHELANKINA, T.S.; KEYLIS-BOROK, V.I.; PAVLOVA, L.G.; REZNYAKOVSKIY, P.T.

Use of electronic computers in locating earthquake epicenters. Dokl. AN SSSR 151 no.2:323-325 J1 '63. (MIRA 16:7)

1. Institut fiziki Zemli im. O.Yu.Shmidta AN SSSR. Predstavleno akademikom Ye.K.Fedorovym.

(Electronic computers) (Seismometry)

APPROVED FOR RELEASE: 09/17/2001 CIA-RDP86-00513R000721620005-8"

ANDRIANOVA, Z.S.; KEYLIS-EOROK, V.I., kand. fiz.-matem. nauk;
LEVSHIN, A.L.; NEYGAUZ, M.G.

[Surface Love waves] Poverkhnostnye volny Liaza. Moskva,
Nauka, 1965. 107 p.

(MIRA 18:4)

KEYLIS-BOROK, V.l., doxto fiz.-matem.rauk

Use of computation mathods in studying the upper manule; symposium in Moscow. Vest. AN SSSR 34 nc.9:88-89 S *64. (MIRA 17:10)

KEYLIS-BOROK, V.I., doktor fiz.-matem.nauk

Computation methods in goophysical research; symposium in Israel. Vest. AN SSSR 35 no.12:54-56 D '65.

(MIRA 19:1)

APPROVED FOR RELEASE: 09/17/2001 CIA-RDP86-00513R000721620005-8"

ACC NR: AT6033690

SOURCE CODE: UR/3231/66/000/002/0003/0045

AUTHOR: Ambol', I.Ya.; Yanovskaya, T.B.; Keylis-Borok, V.I. (Doctor of physico-mathematical sciences

ORG: none

TITLE: Method of combined interpretation of hodographs and amplitude curves in studies of the upper mantle

SOURCE: AN SSSR. Institut fiziki Zemli. Vychislitel'naya seysmologiya, no. 2, 1966. Mashinnaya interpretatsiya seysmicheskikh voln (Machine interpretation of seismic waves), 3-45

TOPIC TAGS: upper mantle, hodograph, seismic wave, Monte Carlo method

计划进行的数据显示。在100

ABSTRACT: The problem of determining the structure of the Earth's upper mantle from the hodographs $t(\Delta)$ of the first few arrivals lacks a unique solution. This ambiguity may be markedly offset, however, if the amplitude curve $A(\Delta)$ or $A^*(\Delta) = \log A(\Delta)$ of direct refracted P waves is also utilized. Travel-time curves which hardly differ in $t(\Delta)$ may be characterized by different $A^*(\Delta)$ and analysis of the latter can result in a marked reduction of the set of curves obtained. The following method of solution of the reciprocal problem is proposed: the sought travel-time curve TTC is parametricized, i.e. represented by a specified

Card 1/2

UDC: 550.34-517:681.142.35

ACC NR. AT6033690

function of the parameters. The possible ranges of variation in these parameters, i.e. the region within which the true travel-time curve exists, are indicated, and individual TTC are scanned, on theoretically calculating for each TTC the data known from observations. The discrepancy between the computed and observational data then is calculated. The set of travel--time curves for which this discrepancy is sufficiently small represents the solution of the problem. Thus, the problem reduces to finding the region of the minimum of multivariate function (discrepancy between calculations and experiment) in the space of unknown parameters of TTC. Finally, the set thus identified must be compactly described, i.e. the common features of all the TTC thus found must be pointed out. The determination of the limits of the parameters and the selection of the type of the parameter-dependent function and the optimal method of search for the minimum depend on the conditions of solution of the converse problem in every particular case: on the accuracy of observational data, type of A (Δ), epicentral spacing for which t (Δ) and A (Δ) are specified, and the nature of the known and unknown parameters. Thus, e.g. if constraints are imposed on both velocities (according to t (Δ)) and velocity gradients (according to A (Δ)), of the methods used to search for the minimum the Monte Carlo method proves to be best. Orig. art. has: 14 figures, 31 formulas, 9 tables.

SUB CODE: 08, - 12/ SUBM DATE: none/ ORIG REF: 012/ OTH REF: 003

Card 2/2

ACC NR: AP7004123

SOURCE CODE: UR/0011/67/000/001/0003/0010

AUTHOR: Keylis-Borok, V. I.

Institute of Geophysics, AN SSSR, Moscow (Institut fiziki Zemli AN SSSR)

TITLE: The use of electronic computers in studying the structure of the earth's crust

SOURCE: AN SSSR. Izvestiya. Seriya geologicheskaya, no. 1, 1967, 3-10

TOPIC TAGS: earth crust, earthquake, electronic computer, computer programming, confured APPLICATION

ABSTRACT: The author presents an attempt to formalize computer operations for seismic data in order to obtain information concerning the structure of the Earth's crust. The first task is to find parameters describing the data succinctly while avoiding the loss of essential information. Earthquakes represent an important characteristic of crustal deformation; they occur in great numbers, each reported by coordinates of the epicenter origin in time, intensity, and so forth. Other parameters are unrelated to tectonics. Total number shocks and energy are more preferable parameters, but still not representative. The author has chosen the sum of energies of individual earthquakes, raised to the 2/3 power, because it is convenient, concise and approximately proportional to the sum of the fracture areas of rupture at the foci. Summation was carried out on

Card 1/2

UDC: 681.142.35:550

ACC NR: APPROVED FOR RELEASE: 09/17/2001 CIA-RDP86-00513R0 a sliding time scale, the sum at any moment being obtained by summing CIA-RDP86-00513R000721620005-8" earthquakes for a certain number of years preceding that moment. This number of years varied for different studies, but was 3, 4, or 5 for examples illustrated in the paper. Consideration was given to areas involved in the summation (such as all the Himalayas for the Assam earthquake, the Pamirs, Hindukush, and Tien Shan for the Khait carthquake). Conclusions concerning the structure of the crust were in conflict because of the problem of selecting data, or the inability to handle all data. The solution of this problem lies in the recognition and elimination of ambiguity. In this study, records of several kinds of waves (longitudinal, transverse, Love, and Rayleigh) were used to investigate a section of the crust between Andizhan and Frunze. Results show a reliability range for seismic velocities down to depths of 60 km on the basis of two models: one with one intracrustal discontinuity (base of the sedimentary layer), the other with two such discontinuities (base of the sedimentary layer and the "granite-basalt" boundary). The : outline of the computer program is broken down into four stages: 1) analysis of records at a single point, 2) study of earthquake foci, 3) refinement of basic empirical patterns, and 4) final conclusions, including sections of the earth, description of seismicity, and methods of recognizing faults. The author concludes that this method of treating seismic data should be more widely used. Orig. art. has: 6 figures. [04]

SUB CODE: 08, 09/ ATD PRESS: 5115 SUBM DATE: 28Apr66/ ORIG REF: 004/ OTH REF: 001/

AND THE PROPERTY AND THE

2/2 Card

Por sj Kraya	spektivy Izucheniya I Ispolizovaniya ya, Goryuchiye Slantsy, 1934, No. 5,				Goryuchi 9.	/ Saratovskogo	
so:	Goryuchiye	Slantey	No. 1931	4-35 TN	.871 . G 74		


KEYL'MAN A. L.

Ispol'zovaniye Zoly Vyurtembergskogo Goryucheqo Slantsa, Goryuchiye Slantsy

1934, No. 6,45.

SO: Gory uchiye Slantey #1934-35 TN. 871 674

Combined solubility table of electrolytes. Khim.v shkole 11 no.5:71-72 S-0 *56. (MLRA 9:11) (Electrolytes-Tables) (Solubility-Tables)



KEYL'MAN, G.A.

New data on the geological structure of the Sysert' granitic region [with summary in English]. Sov. geol. 1 no.6:121-135 '58. (MIRA 11:10)

1. Ural'skoye geologicheskoye upravleniye. (Ural Mountains--Geology)

KEYL'MAN, G.A.


Geology of the Ufaley granite massif. Sov. geol. 6 no.10: 120-122 0 '63. (MIRA 17:1)

1. Ural'skoye geologicheskoye upravleniye.

APPROVED FOR RELEASE: 09/17/2001 CIA-RDP86-00513R000721620005-8"

A PART OF THE PART

KEYL MAN, V. A.: Haster Tech Sci (diss) -- "Methods of improving the dirt roads on loss in Rostov Oblast". Moscow, 1959. 25 pp (Min Higher Educ USSR, Moscow Automobile and Road Inst), 150 copies (KL, No 13, 1959, 105)



28194 S/194/61/000/005/008/078 D201/D303

9.9865 3,9300 (1019,1109,1327)

Gol'tsman, F.M. and Keyl'man, Yu.N.

AUTHORS:

A universal filter for seismic signals

TITLE:

PERIODICAL:

Referativnyy zhurnal. Avtomatika i radioelektronika. no. 5, 1961, 21-22, abstract 5 Al61 (V sb. Prikl. geofizika, no. 25, M., 1960, 55-56)

The theory is briefly explained of a universal filter with any pre-determined frequency response. A description is given of a filter in which an audio amplifier in the form of a delay line is TEXT: This makes it possible to dispense with a preliminary recording and subsequent reproduction of signals being filtered. total delay time of the signal in 120 LC sections of the lelay line is 72 microseconds, frequency range 10-150 c/s. The amplitude response of amplifiers is linear for the range of input voltages 0-0.1 V. The output of the instrument was designed for connection to a normal loop oscillograph, The technique of using the instru-

Card 1/2

APPROVED FOR RELEASE: 09/17/2001 CIA-RDP86-00513R000721620005-8"

enne comunicate morneciae comunicati

28194

S/194/61/000/005/008/078 D201/D303

A universal filter for seismic signals

ment is described and results of testing it are given. bility is emphasized of using the instrument in seismological investigations, in particular for frequency separation of magnetic and gravitational anomalies. 7 figures. 5 references. / Abstracter's note: Complete translation

KEYL'MAN, Yu.N.

An attempt of model studies on self-reproductive and evolutionary processes. Trudy Len. ob-va est. 72 no.1:134-137 '61. (MIRA 15:3)

(INFORMATION THEORY IN BIOLOGY)

KEYLKE, L. I.

Draina_ee

Hardre-Fillber ap gradum for lowering water level. Seret. prom. 42, no. 1, 1951

Monthly List of Russian Accessions, Library of Congress, August 1952. Unclassified.

KUTHARR, L. J.

Water, Unlerground

Needle-Filter asparatus for lowering mater level. Stroi. pros. 29, no. 12, 1951

Monthly List of Russian Accessions, Library of Congress, August 1952. Unclassified.

ETHATE, L. I.

Univariatembs

Overall machanization of soil operations in erecting ambanks anto. Stroi. prom., 30, No. 7, 1952.

Monthly Lieb of Russian Accessions, Library of Congress, October 1954. Unclassified.

- 1. KETMAKH, L.I. Eng.
- 2. USSR (600)
- 4. Chimneys
- Technical rules for building a reinforced concrete chimney. Stroi.prom. 30 no. 10 0 *52.

9. Monthly List of Russian Accessions, Library of Congress, January 1953, Unclassified.